

Perturbed threat monitoring following a traumatic event predicts risk for post-traumatic stress disorder

R. Naim^{1*}, I. Wald¹, A. Lior², D. S. Pine³, N. A. Fox⁴, G. Sheppes¹, P. Halpern^{2,5} and Y. Bar-Haim^{1,6}

¹School of Psychological Sciences, Tel Aviv University, Israel

²Emergency Medicine Department, Tel Aviv Medical Center, Israel

³National Institute of Mental Health, Bethesda, MD, USA

⁴Department of Human Development, University of Maryland, College Park, MD, USA

⁵Sackler Faculty of Medicine, Tel Aviv University, Israel

⁶The Sagol School of Neuroscience, Tel Aviv University, Israel

Background. Post-traumatic stress disorder (PTSD) is a chronic and difficult to treat psychiatric disorder. Objective, performance-based diagnostic markers that uniquely index risk for PTSD above and beyond subjective self-report markers could inform attempts to improve prevention and early intervention. We evaluated the predictive value of threat-related attention bias measured immediately after a potentially traumatic event, as a risk marker for PTSD at a 3-month follow-up. We measured the predictive contribution of attentional threat bias above and beyond that of the more established marker of risk for PTSD, self-reported psychological dissociation.

Method. Dissociation symptoms and threat-related attention bias were measured in 577 motor vehicle accident (MVA) survivors (mean age=35.02 years, 356 males) within 24 h of admission to an emergency department (ED) of a large urban hospital. PTSD symptoms were assessed at a 3-month follow-up using the Clinician-Administered PTSD Scale (CAPS).

Results. Self-reported dissociation symptoms significantly accounted for 16% of the variance in PTSD at follow-up, and attention bias toward threat significantly accounted for an additional 4% of the variance in PTSD.

Conclusions. Threat-related attention bias can be reliably measured in the context of a hospital ED and significantly predicts risk for later PTSD. Possible mechanisms underlying the association between threat bias following a potentially traumatic event and risk for PTSD are discussed. The potential application of an attention bias modification treatment (ABMT) tailored to reduce risk for PTSD is suggested.

Received 9 May 2013; Revised 22 August 2013; Accepted 31 August 2013; First published online 17 October 2013

Key words: Anxiety, attention bias, dissociation, PTSD.

Introduction

Following trauma, most people experience stress-related symptoms but these typically resolve over a relatively short period of time. An important minority who continue to experience post-traumatic stress symptoms after 4 weeks are diagnosed with post-traumatic stress disorder (PTSD) (Rothbaum *et al.* 1992; Blanchard *et al.* 1995; Riggs *et al.* 1995; Koren *et al.* 1999). Diagnostic markers that identify this unique subgroup could inform attempts to improve early intervention and prevention (Bryant *et al.* 1999, 2008). In this study we focused on the predictive risk of two cognitive markers measured immediately following a potentially traumatic event: self-reported symptoms of psychological dissociation and behaviorally measured threat-related attention bias.

Monitoring potential threats in the environment is essential for survival. The human threat-monitoring system relies on a dedicated neural network that continually balances the cognitive resources allocated to the evaluation of potential threats and coordinates ongoing psychological and behavioral response patterns (Liddell *et al.* 2005; Pessoa & Adolphs, 2010; Adolphs, 2013). Traumatic events can challenge the delicate cognitive-behavioral balance maintained by the threat-monitoring system, and could manifest as threat-related cognitive biases (Ehlers & Clark, 2000). Importantly, such perturbations in the threat-monitoring system seem to resemble two primary symptom clusters of PTSD, namely threat avoidance/dissociation and hypervigilance (Scully, 2000).

High levels of peritraumatic dissociation symptoms have been identified as a potential risk marker for PTSD, with meta-analyses reporting a medium effect size (Ozer *et al.* 2003; Lensvelt-Mulders *et al.* 2008). Symptoms of psychological dissociation during and immediately following a traumatic experience are

* Address for correspondence: R. Naim, MA, School of Psychological Sciences, Tel Aviv University, Tel Aviv 69978, Israel.
(Email: reutnaim@post.tau.ac.il)

fairly common (Bremner *et al.* 1992; Koopman *et al.* 1994; Marmar *et al.* 1994, 1999) and involve reduction in awareness to one's surroundings and altered perceptions of self and the environment (Cardena & Spiegel, 1993). Although dissociation symptoms could be adaptive in the short run by allowing traumatic perceptions and experiences to be split from awareness and thus reduce immediate distress, they also index later risk for PTSD (van der Kolk *et al.* 1996; Bremner & Brett, 1997; Harvey & Bryant, 1998, 2002; Marmar *et al.* 1999; Briere *et al.* 2005).

Another potential cognitive marker of risk for PTSD following trauma emerges from the research on threat-related attention biases in anxiety. Attention bias toward threats represents one of the most consistently demonstrated correlates of anxiety disorders including PTSD (Yiend, 2010). The attention of patients with PTSD is selectively biased toward threat-related information (Thrasher *et al.* 1994; Litz *et al.* 1996; Bryant & Harvey, 1997; Moradi *et al.* 1999; Jenkins *et al.* 2000). However, these findings relate to patients with ongoing PTSD and do not reveal whether peri-traumatic attention bias is predictive of PTSD risk. One study tested longitudinally whether attention bias toward threat following a traumatic event predicted later risk for PTSD, and found null results (Elsesser *et al.* 2005). In that study participants from a range of potentially traumatic exposures were included, the task used to measure attention bias used complex pictures depicting neutral and traumatic events, and attention bias was first assessed approximately 1 month after the trauma, with large variability in post-event testing times (7–56 days).

In the current study, to reduce potential experimental variability, we tested whether threat-related attention bias measured within 24 h after a specific traumatic event (motor vehicle accident, MVA) and using visually less complex standardized word stimuli may serve as a predictive marker of PTSD at the 3-month follow-up. We expected that threat vigilance would confer higher risk for PTSD above and beyond the established predictive value of self-reported dissociation symptoms. Direct measurement of perturbations in the threat-monitoring system using computerized tasks could highlight aspects of risk that are not accessible from self-reports on symptoms. Such direct measurement also bypasses the concern of shared method variance between patient-reported dissociation symptoms and patient-reported PTSD symptoms as outcome.

Method

Participants

MVA survivors with minor injuries, who were admitted to the emergency department (ED) at the

Soraski Medical Center, Tel-Aviv between March 2011 and January 2012, formed the potential pool of participants in the study. Data were collected 5 days a week in two 5-h shifts/day. During these shifts, every potential participant was approached for recruitment. Inclusion criteria were: (a) being involved in an MVA and fulfilling DSM-IV criterion A1 of the PTSD diagnosis; (b) suffering only minor injuries as determined by the ED medical staff; specifically, subjects could not suffer internal bleeding, head injury or require urgent surgery; (c) age between 18 and 65 years; (d) fluent in Hebrew; and (e) the MVA had occurred within the past 24 h.

Of the 618 potentially eligible individuals, 41 declined participation, generating data in 577 (mean age=35.02 years, s.d.=11.68, range 18–65, 356 males). Of these 577, 194 were drivers of four-wheel vehicles, 248 were motorcyclists, 42 were bicycle riders, 30 were pedestrians, and 63 were passengers. Three months later, an attempt was made to contact these 577, among whom 24 could not be contacted and 138 declined further participation. This left 415 participants with full data, representing 71.9% of the initial sample. Participants with full data sets did not differ from other subjects in terms of most variables acquired at baseline. These included age, gender distribution, threat-related attention bias, and most types of symptoms (all p 's > 0.17). However, relative to participants with full data, non-participants had more dissociation symptoms following the MVA (6.2 *v.* 9.3, $t_{406}=2.46$, $p=0.014$).

Written informed consent was obtained from all participants. Given the sensitivity of research in the ED setting, research staff ensured that participants clearly understood the voluntary nature of the study. The study was approved by the Tel Aviv University Institutional Review Board and the Ethics Committee of the Sourasky Medical Center.

Clinician-Administered PTSD Scale (CAPS)

PTSD was diagnosed using the CAPS and served as the outcome measure of the study. The CAPS is a structured interview used to make current or lifetime diagnosis of PTSD. The first 17 items of the CAPS correspond to the DSM-IV criteria for PTSD (Blake *et al.* 1995). For each item, symptom intensity (from 0=no distress to 4=severe distress) and frequency (from 0=never to 4=almost every day) over the past month are evaluated. The impact of symptoms on social and occupational functioning and overall PTSD severity are also measured. The three symptom clusters of PTSD are indexed: intrusive recollections (cluster B), avoidant/numbing symptoms (cluster C) and hyperarousal (cluster D). A symptom is considered present

if it has a frequency of ≥ 1 and an intensity of ≥ 2 . A PTSD diagnosis required at least one B-cluster symptom, three C-cluster symptoms and two D-cluster symptoms, in addition to meeting the other diagnostic criteria of at least 1 month's duration and significant distress or impairment in functioning. CAPS total score was derived by summing the first 17 items. The CAPS is considered the gold standard in PTSD diagnosis and possesses good sensitivity, specificity, retest reliability and validity (Blake *et al.* 1995; Blanchard *et al.* 1996). Interviewers were four graduate clinical psychology students trained and supervised to 85% reliability criterion with an experienced clinical psychologist. Internal consistency in the current study was good (Cronbach's $\alpha=0.95$).

Primary predictors of PTSD

Clinician-Administered Dissociative States Scale (CADSS)

The CADSS (Bremner *et al.* 1998) is designed to assess the occurrence and severity of dissociative symptoms in relation to the participant's perceptual experience during the interview. The items assess how perceptually 'in touch' (or 'out of touch') an individual is with the environment following exposure to stressful conditions. The scale contains 19 items rated on a Likert-type scale from 0 (not at all) to 4 (extremely). Cronbach's α in the current study was 0.75.

Threat bias assessment: the dot-probe task

The dot-probe task used in the current study was similar to the one used by Bar-Haim *et al.* (2010) and Wald *et al.* (2011a). In brief, 25 word pairs, each consisting of one general threat word (e.g. scared, danger, dead) and one neutral word (e.g. line, tomatoes, carpets), were used as stimuli. Words of each pair were matched on number of letters and frequency of usage in Hebrew. Each trial began with a 500-ms fixation display '+++' in the center of the screen followed by a word pair presented in white block text 1-cm high. One word appeared directly above and the other directly below the location vacated by the preceding fixation signal. A distance of 1.5 cm separated the two words. The word pair remained on the screen for 1000 ms, and was then replaced by a target probe (either '..' or '..') that appeared in one of the two locations vacated by the words[†]. Participants had to identify the probe types. The participant's response cleared the screen and the next trial began 500 ms later.

The task comprised 100 trials and took approximately 4 min to complete. Probes appeared with equal frequency at the location of either the threat or

the neutral word. The speed with which participants discriminated the probe type accurately was recorded on each trial. Trials with an incorrect response and trials in which the response time was more than 2 standard deviations of the participant's mean for a particular condition were excluded from subsequent analysis (<2% of all trials). Threat bias was calculated as the difference between the average response time to targets appearing at neutral word locations and those appearing at threat word locations. Positive bias values represent attention bias toward threat and negative values reflect an attentional bias away from threat. Word valence location, target location and target type were fully counterbalanced.

Baseline PTSD and depression symptoms

Nine-item Patient Health Questionnaire (PHQ-9)

The PHQ-9 (Spitzer *et al.* 1999) is a self-reported depression rating scale consisting of nine items on which the DSM-IV diagnosis of major depressive disorder is based. Each item is scored on a Likert-type scale ranging from 0 (not at all) to 3 (nearly every day). A total score is obtained by summing across all items. An additional item assesses the severity of impairment. Reliability and diagnostic validity of the PHQ-9 have been established in several studies (Spitzer *et al.* 1999; Kroenke *et al.* 2001; Kroenke & Spitzer, 2002). Cronbach's α in the current study was 0.78.

PTSD Checklist (PCL)

The prevalence of PTSD symptoms before the MVA was evaluated using the 17-item National Center for PTSD Checklist of the Department of Veterans Affairs, the PCL (Blanchard *et al.* 1996; Kang *et al.* 2003; Hoge *et al.* 2004). Symptoms were related to any stressful experience (in the wording of the 'specific stressor' version of the checklist). Scores can range from 17 to 85, with higher scores reflecting more symptoms of PTSD. Cronbach's α in the current study was 0.92.

Procedure

Participants were assessed twice 3 months apart. In the first assessment, at the ED within 24 h of their MVA, participants completed the CADSS, the dot-probe task, the PCL and the PHQ-9. At the second assessment, 3 months later, participants were diagnosed for PTSD in a telephone interview using the CAPS.

Data analysis

Hierarchical multivariate logistic regression analysis was conducted to examine the predictive contribution

[†] The notes appear after the main text.

Table 1. Background characteristics and predictor measures within 24 h of the MVA by PTSD diagnosis at the 3-month follow-up

	PTSD	No PTSD	<i>p</i> value
Gender (%)			
Male	4.5	95.5	0.01
Female	10.8	89.2	
Family status (%)			
Single	12.5	87.5	0.73
Married	8.1	91.9	
Participant's role in the MVA (%)			
Driver	5.2	94.8	0.002
Passenger	17.0	83.0	
Age (years), mean (s.d.)	34.75 (11.54)	33.71 (11.16)	0.64
PCL (baseline PTSD), mean (s.d.)	32.03 (15.59)	24.42 (10.24)	0.02
PHQ-9 (depression), mean (s.d.)	7.77 (6.79)	5.15 (5.04)	0.05
Attention threat bias (ms), mean (s.d.)	23 (45)	1 (33)	0.02
Dissociation, mean (s.d.)	22.77 (13.31)	5.29 (7.77)	0.0001

MVA, Motor vehicle accident; PTSD, post-traumatic stress disorder; PCL, PTSD Checklist; PHQ-9, nine-item Patient Health Questionnaire; s.d.; standard deviation.

of dissociation symptoms and threat-related attention bias collected within 24 h of an MVA to vulnerability to PTSD at the 3-month follow-up. Baseline self-reported PTSD and depression symptoms (PCL and PHQ-9) were entered into the regression model first. Because PTSD at follow-up was more prevalent in females than males and in passengers than drivers, gender and role in the MVA were also entered into the regression in the first step. The primary predictors (dissociation symptoms and threat-related attention bias) were entered into the model in the second step. The interaction between dissociation and attention bias was entered in the third step.

Results

Descriptive statistics for all variables are provided in Table 1. Zero-order correlations between the different measures collected at baseline are presented in Table 2. Of the 415 participants who provided data at follow-up, 28 (6.75%) had PTSD based on the CAPS. Participants diagnosed with PTSD did not differ from the non-PTSD group in age or gender distribution (p 's > 0.10). Nevertheless, the two groups differed in pre-MVA self-reported PTSD symptoms (PCL) and depression symptoms (PHQ-9) as measured immediately after the MVA ($t_{s413}=2.54$ and 2.0 , p 's = 0.02 and 0.055 respectively), with higher symptoms in the PTSD than the non-PTSD group. Within the group with PTSD at follow-up, there were more females than males (10.8% *v.* 4.5%, $\chi^2=6.04$, $p=0.014$) and

Table 2. Zero-order correlations between baseline measures of PTSD (PCL), depression (PHQ-9), dissociation (CADSS), and attentional threat bias and CAPS total scores

	Threat bias	PCL	PHQ-9	CADSS
PCL	0.09*			
PHQ-9	-0.03	0.63***		
CADSS	0.10**	0.48***	0.38***	
CAPS	0.12**	0.22***	0.19***	0.55***

PTSD, Post-traumatic stress disorder; PCL, PTSD Checklist; PHQ-9, nine-item Patient Health Questionnaire; CADSS, Clinician-Administered Dissociative States Scale; CAPS, Clinician-Administered PTSD Scale.

* $p=0.052$, ** $p<0.05$, *** $p<0.0001$.

more passengers than drivers (17.0% *v.* 5.2%, $\chi^2=9.59$, $p=0.008$).

The regression model accounted for 38% of the variance in PTSD diagnosis at follow-up ($\chi^2=42.8$, $p<0.001$). The four background variables entered in step 1 (PCL, PHQ-9, gender, and role in the MVA) significantly accounted for 18% of the variance in PTSD at follow-up ($\chi^2=20.32$, $p<0.001$). Step 2 of the regression revealed that higher levels of dissociation immediately following the MVA were associated with greater risk for PTSD at the 3-month follow-up [$B=0.09$, odds ratio (OR) 1.10, 95% confidence interval (CI) 1.05–1.15, Wald=16.10, $p<0.0001$], accounting for an additional 16% of the variance in PTSD diagnosis. Greater attention bias toward threat at baseline was

Table 3. Estimated coefficients, standard errors and 95% CIs for predictors in the three steps of the regression model predicting PTSD

	Predictor	B	s.e.	Wald	Exp (B)	95% CI
Step 1	Baseline PCL	0.02	0.03	0.51	1.02	0.96–1.08
	Role in the MVA	0.50	0.50	1.00	1.66	0.62–4.46
	Baseline PHQ-9	0.03	0.07	0.17	0.97	0.85–1.11
Step 2	Gender	1.20	0.69	3.08	3.35	0.87–12.90
	Dissociation	0.09**	0.023	16.10	1.10	1.05–1.15
Step 3	Attention bias	0.02*	0.008	4.50	1.02	1.001–1.03
	Dissociation-by-attention bias	0.001	0.001	1.39	1.00	1.00–1.00

PTSD, Post-traumatic stress disorder; PCL, PTSD Checklist; MVA, motor vehicle accident; PHQ-9, nine-item Patient Health Questionnaire; B, unstandardized estimated coefficient; s.e., standard error; CI, confidence interval.

* $p < 0.05$, ** $p < 0.0001$.

also associated with higher incidence of PTSD at follow-up, accounting for an additional 4% of the variance above and beyond the variance accounted for by the variables entered in step 1 and the dissociation symptoms ($B=0.02$, OR 1.02, 95% CI 1.001–1.03, Wald=4.50, $p < 0.04$). Finally, the interaction term between dissociation and attention bias did not predict PTSD diagnosis ($\chi^2=1.53$, $p > 0.20$). The estimate coefficients for the regression model are shown in Table 3.

Discussion

The novel finding of the current study is that performance-based attention bias toward threat, measured immediately after an MVA, predicted PTSD above and beyond the variability predicted by self-reported dissociation symptoms, which was previously found to be a strong predictor of risk for PTSD (Lensvelt-Mulders *et al.* 2008), and specifically in MVA survivors (Ehlers *et al.* 1998; Holeva & Tarrier, 2001).

One possible mechanism linking enhanced threat-related attention bias to risk for later PTSD relates to greater accessibility and enhanced consolidation of trauma-related elements that is facilitated by increased threat attendance. Increased attention to the traumatic event could extend the range of accessible trauma-related triggers and thereby increase the risk for PTSD symptoms. A mechanism that may operate in a similar way but in a different system has been suggested in studies demonstrating that enhanced priming for trauma-related stimuli measured shortly after the traumatic event predicted subsequent flashbacks (Michael *et al.* 2005), or re-experiencing of the trauma (Ehlers *et al.* 2006). More research is needed to establish the validity of such mechanisms in relation to attentional threat bias and enhanced PTSD symptoms.

Enhanced low-level attention toward threats following a traumatic event could also enhance the psychological tendency to dissociate. Specifically, hyper-engagement with threats at the basic attention allocation level could feed forward to induce intolerable stress that, in some people, invokes psychological defenses in the form of dissociative symptoms. Indeed, in the current sample, a small but significant association was found between threat bias and dissociation symptoms. Similar vigilance-avoidance models of attentional engagement have been proposed for other anxiety disorders in which dissociation is not a typical symptom (Mogg *et al.* 2004; Koster *et al.* 2005; Pflugshaupt *et al.* 2005). It may be the case that the high intensity of threat associated with traumatic events causes more minor patterns of attentional avoidance to propagate to other cognitive systems and manifest more robustly in the form of dissociation. Evidence from neuroimaging research in PTSD patients lends support this possibility of a biphasic process of initial attentional vigilance toward threats and later threat avoidance (Adenauer *et al.* 2010).

The results of the current study should also be discussed in relation to prior reports of plasticity in threat-related attention bias under conditions of acute stress. Unlike the typical finding of threat vigilance in anxious participants under neutral laboratory conditions (Bar-Haim *et al.* 2007), several studies have shown that, in normative samples, imminent threat can produce attentional threat avoidance rather than vigilance (Beevers *et al.* 2011; Wald *et al.* 2011a; Shechner *et al.* 2012). Such threat avoidance under stress has been associated with risk for PTSD symptoms (Wald *et al.* 2011b). Thus, why, in the context of the current study, does threat vigilance rather than threat avoidance conferred risk for later PTSD? The answer to this question may relate to the different contexts in which these previous studies were conducted

relative to the context of the current study. Studies that found attentional threat avoidance to predict PTSD symptoms were conducted under conditions of immediate and imminent threat (e.g. ongoing rocket attack, threat of an electric shock, or military deployment). In these situations an adaptive response involves attending to threats, making attentional threat avoidance a maladaptive response that has been associated with risk of PTSD symptoms. By contrast, the minor severity of injuries of participants in the current study, and the timing and location of attention bias measurement relative to the trauma (threat was no longer present and the participants were in the caring environment of the ED), may have rendered the situation relatively safe. Therefore, the adaptive response would be to ignore minor threats, and a maladaptive association between hypervigilance for minor threats and anxiety-related symptoms emerged as a risk factor for PTSD. Additional studies are needed to further clarify the interplay between threat-related attention patterns and context and their relationship to PTSD.

The current study also has limitations that future research may overcome. First, although the results indicate that threat-related attention bias and dissociation symptoms predict increased risk for PTSD, they do not provide information about participants' threat-related attention or dissociative tendencies prior to the traumatic event. Therefore, the correlation between these predictors and PTSD could be driven either by pre-trauma cognitive styles acting as personal vulnerability factors or by differential plasticity in participants' cognitive reactions to the traumatic event. The resolution of this issue may point to threat bias and dissociation as general risk factors for anxiety and stress-related psychopathology (Bar-Haim *et al.* 2007), or as indices more closely associated with responses to traumatic exposure. Second, in the current study threat-related attention bias and dissociation were not measured at follow-up and thus do not allow inference about the potential role of persistent threat bias and dissociation in PTSD development. Future studies measuring threat-related attention bias and dissociation at various time points before and after exposure to traumatic events are needed to clarify these conceptual issues (Wald *et al.* 2013).

In conclusion, the current findings contribute an additional piece to the puzzle of PTSD vulnerability factors that can be measured efficiently and at a low cost soon after the occurrence of traumatic events. Nevertheless, the combined percentage of the variance in risk for PTSD accounted for by the current predictors is relatively small (38%), and thus renders their clinical utility unclear. Additional research is needed to establish the role of threat-related attention bias in

vulnerability to PTSD and its interaction with other factors that could reliably predict PTSD development. Identification of the risk associated with enhanced vigilance toward threats in the ED could lead to the development of computer-based interventions designed to modify such attention biases, with the intention of reducing risk for PTSD (Pine *et al.* 2009). Attention bias modification treatments (ABMTs) have shown efficacy in the treatment of anxiety disorders (Bar-Haim, 2010; Hakamata *et al.* 2010). The current results suggest that there may also be a potential for an ABMT as a prevention protocol for those at increased risk for PTSD following traumatic exposures (See *et al.* 2009).

Note

- ¹ The selection of the 1000-ms presentation for the word stimuli was based on the fact that, across traditional supraliminal assessment implementations of the dot-probe task, the stimulus onset asynchrony (SOA) between words and probes has ranged from 28 ms (e.g. Mogg & Bradley, 1999) to 1000 ms (e.g. Hunt *et al.* 2007; Clarke *et al.* 2008), or occasionally even longer (e.g. Shane & Peterson, 2007; for a review, see Bar-Haim *et al.* 2007). Selection of the upper end of the commonly used SOA range was driven by an attempt to maximize the prospect of participants cognitively registering, and attentionally responding to, the word stimuli. In addition, our own experience has been successful with using the 1000-ms version of the task in four different studies related to traumatic exposure in normative populations (see Bar-Haim *et al.* 2010; Wald *et al.* 2011a,b; Shechner *et al.* 2012). Thus, to facilitate comparison with these previous studies we opted to maintain the 1000-ms presentation.

Acknowledgments

This work was supported by the Israel Insurance Association, Association of Life Insurance Companies of Israel (0610714692). Professor Y. Bar-Haim is independent of any commercial funder and had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Declaration of Interest

None.

References

- Adenauer H, Pinosch S, Catani C, Gola H, Keil J, Kissler J, Neuner F (2010). Early processing of threat cues in posttraumatic stress disorder – evidence for a cortical

- vigilance-avoidance reaction. *Biological Psychiatry* **68**, 451–458.
- Adolphs R** (2013). The biology of fear. *Current Biology* **23**, R79–R93.
- Bar-Haim Y** (2010). Research review: attention bias modification (ABM): a novel treatment for anxiety disorders. *Journal of Child Psychology and Psychiatry* **51**, 859–870.
- Bar-Haim Y, Holoshitz Y, Eldar S, Frenkel TI, Muller D, Charney DS, Pine DS, Fox NA, Wald I** (2010). Life-threatening danger and suppression of attention bias to threat. *American Journal of Psychiatry* **167**, 694–698.
- Bar-Haim Y, Lamy D, Pergamin L, Bakermans-Kranenburg MJ, van IJzendoorn MH** (2007). Threat-related attentional bias in anxious and non-anxious individuals: a meta-analytic study. *Psychological Bulletin* **133**, 1–24.
- Beavers CG, Lee HJ, Wells TT, Ellis AJ, Telch MJ** (2011). Association of predeployment gaze bias for emotion stimuli with later symptoms of PTSD and depression in soldiers deployed in Iraq. *American Journal of Psychiatry* **168**, 735–741.
- Blake DD, Weathers FW, Nagy LM, Kaloupek DG, Gusman FD, Charney DS, Keane TM** (1995). The development of a Clinician-Administered PTSD Scale. *Journal of Traumatic Stress* **8**, 75–90.
- Blanchard EB, Hickling EJ, Mitnick N, Taylor AE, Loos WR, Buckley TC** (1995). The impact of severity of physical injury and perception of life threat in the development of post-traumatic stress disorder in motor vehicle accident victims. *Behaviour Research and Therapy* **33**, 529–534.
- Blanchard EB, Jones-Alexander J, Buckley TC, Forneris CA** (1996). Psychometric properties of the PTSD Checklist (PCL). *Behaviour Research and Therapy* **34**, 669–673.
- Bremner JD, Brett E** (1997). Trauma-related dissociative states and long-term psychopathology in posttraumatic stress disorder. *Journal of Traumatic Stress* **10**, 37–49.
- Bremner JD, Krystal JH, Putnam FW, Southwick SM, Marmar C, Charney DS, Mazure CM** (1998). Measurement of dissociative states with the Clinician-Administered Dissociative States Scale (CADSS). *Journal of Traumatic Stress* **11**, 125–136.
- Bremner JD, Southwick S, Brett E, Fontana A, Rosenheck R, Charney DS** (1992). Dissociation and posttraumatic stress disorder in Vietnam combat veterans. *American Journal of Psychiatry* **149**, 328–332.
- Briere J, Scott C, Weathers F** (2005). Peritraumatic and persistent dissociation in the presumed etiology of PTSD. *American Journal of Psychiatry* **162**, 2295–2301.
- Bryant RA, Harvey AG** (1997). Attentional bias in posttraumatic stress disorder. *Journal of Traumatic Stress* **10**, 635–644.
- Bryant RA, Mastrodomenico J, Felmingham KL, Hopwood S, Kenny L, Kandris E, Cahill C, Creamer M** (2008). Treatment of acute stress disorder: a randomized controlled trial. *Archives of General Psychiatry* **65**, 659–667.
- Bryant RA, Sackville T, Dang ST, Moulds M, Guthrie R** (1999). Treating acute stress disorder: an evaluation of cognitive behavior therapy and supportive counseling techniques. *American Journal of Psychiatry* **156**, 1780–1786.
- Cardena E, Spiegel D** (1993). Dissociative reactions to the San Francisco Bay Area earthquake of 1989. *American Journal of Psychiatry* **150**, 474–478.
- Clarke P, MacLeod C, Shirazee N** (2008). Prepared for the worst: readiness to acquire threat bias and susceptibility to elevate trait anxiety. *Emotion* **8**, 47–57.
- Ehlers A, Clark DM** (2000). A cognitive model of posttraumatic stress disorder. *Behaviour Research and Therapy* **38**, 319–345.
- Ehlers A, Mayou RA, Bryant B** (1998). Psychological predictors of chronic posttraumatic stress disorder after motor vehicle accidents. *Journal of Abnormal Psychology* **107**, 508–519.
- Ehlers A, Michael T, Chen YP, Payne E, Shan S** (2006). Enhanced perceptual priming for neutral stimuli in a traumatic context: a pathway to intrusive memories? *Memory* **14**, 316–328.
- Elsesser K, Sartory G, Tackenberg A** (2005). Initial symptoms and reactions to trauma-related stimuli and the development of posttraumatic stress disorder. *Depression and Anxiety* **21**, 61–70.
- Hakamata Y, Lissek S, Bar-Haim Y, Britton JC, Fox NA, Leibenluft E, Ernst M, Pine DS** (2010). Attention bias modification treatment: a meta-analysis toward the establishment of novel treatment for anxiety. *Biological Psychiatry* **68**, 982–990.
- Harvey AG, Bryant RA** (1998). The effect of attempted thought suppression in acute stress disorder. *Behaviour Research and Therapy* **36**, 583–590.
- Harvey AG, Bryant RA** (2002). Acute stress disorder: a synthesis and critique. *Psychological Bulletin* **128**, 886–902.
- Hoge CW, Castro CA, Messer SC, McGurk D, Cotting DI, Koffman RL** (2004). Combat duty in Iraq and Afghanistan, mental health problems, and barriers to care. *New England Journal of Medicine* **351**, 13–22.
- Holeva V, Tarrier N** (2001). Personality and peritraumatic dissociation in the prediction of PTSD in victims of road traffic accidents. *Journal of Psychosomatic Research* **51**, 687–692.
- Hunt C, Keogh E, French CC** (2007). Anxiety sensitivity, conscious awareness and selective attentional biases in children. *Behaviour Research and Therapy* **45**, 497–509.
- Jenkins MA, Langlais PJ, Delis D, Cohen RA** (2000). Attentional dysfunction associated with posttraumatic stress disorder among rape survivors. *Clinical Neuropsychologist* **14**, 7–12.
- Kang HK, Natelson BH, Mahan CM, Lee KY, Murphy FM** (2003). Post-traumatic stress disorder and chronic fatigue syndrome-like illness among Gulf War veterans: a population-based survey of 30000 veterans. *American Journal of Epidemiology* **157**, 141–148.
- Koopman C, Classen C, Spiegel D** (1994). Predictors of posttraumatic stress symptoms among survivors of the Oakland/Berkeley, Calif., firestorm. *American Journal of Psychiatry* **151**, 888–894.
- Koren D, Arnon I, Klein E** (1999). Acute stress response and posttraumatic stress disorder in traffic accident victims: a one-year prospective, follow-up study. *American Journal of Psychiatry* **156**, 367–373.

- Koster EHW, Verschuere B, Crombez G, Van Damme S** (2005). Time-course of attention for threatening pictures in high and low trait anxiety. *Behaviour Research and Therapy* **43**, 1087–1098.
- Kroenke K, Spitzer RL** (2002). The PHQ-9: a new depression diagnostic and severity measure. *Psychiatric Annals* **32**, 509–515.
- Kroenke K, Spitzer RL, Williams JBW** (2001). The PHQ-9: validity of a brief depression severity measure. *Journal of General Internal Medicine* **16**, 606–613.
- Lensvelt-Mulders G, van der Hart O, van Ochten JM, van Son MJM, Steele K, Breeman L** (2008). Relations among peritraumatic dissociation and posttraumatic stress: a meta-analysis. *Clinical Psychology Review* **28**, 1138–1151.
- Liddell BJ, Brown KJ, Kemp AH, Barton MJ, Das P, Peduto A, Gordon E, Williams LM** (2005). A direct brainstem-amygdala-cortical ‘alarm’ system for subliminal signals of fear. *NeuroImage* **24**, 235–243.
- Litz BT, Weathers FW, Monaco V, Herman DS, Wulfsohn M, Marx B, Keane TM** (1996). Attention, arousal, and memory in posttraumatic stress disorder. *Journal of Traumatic Stress* **9**, 497–519.
- Marmar CR, Weiss DS, Metzler TJ, Delucchi KL, Best SR, Wentworth KA** (1999). Longitudinal course and predictors of continuing distress following critical incident exposure in emergency services personnel. *Journal of Nervous and Mental Disease* **187**, 15–22.
- Marmar CR, Weiss DS, Schlenger WE, Fairbank JA, Jordan BK, Kulka RA, Hough RL** (1994). Peritraumatic dissociation and posttraumatic stress in male Vietnam theater veterans. *American Journal of Psychiatry* **151**, 902–907.
- Michael T, Ehlers A, Halligan SL** (2005). Enhanced priming for trauma-related material in posttraumatic stress disorder. *Emotion* **5**, 103–112.
- Mogg K, Bradley BP** (1999). Orienting of attention to threatening facial expressions presented under conditions of restricted awareness. *Cognition and Emotion* **13**, 713–740.
- Mogg K, Bradley BP, Miles F, Dixon R** (2004). Time course of attentional bias for threat scenes: testing the vigilance-avoidance hypothesis. *Cognition and Emotion* **18**, 689–700.
- Moradi AR, Taghavi MR, Doost HTN, Yule W, Dalgleish T** (1999). Performance of children and adolescents with PTSD on the Stroop colour-naming task. *Psychological Medicine* **29**, 415–419.
- Ozer EJ, Best SR, Lipsey TL, Weiss DS** (2003). Predictors of posttraumatic stress disorder and symptoms in adults: a meta-analysis. *Psychological Bulletin* **129**, 52–73.
- Pessoa L, Adolphs R** (2010). Emotion processing and the amygdala: from a ‘low road’ to ‘many roads’ of evaluating biological significance. *Nature Reviews Neuroscience* **11**, 773–782.
- Pflugshaupt T, Mosimann UP, von Wartburg R, Schmitt W, Nyffeler T, Muri RM** (2005). Hypervigilance-avoidance pattern in spider phobia. *Journal of Anxiety Disorders* **19**, 105–116.
- Pine DS, Helfinstein SM, Bar-Haim Y, Nelson E, Fox NA** (2009). Challenges in developing novel treatments for childhood disorders: lessons from research on anxiety. *Neuropsychopharmacology* **34**, 213–228.
- Riggs DS, Rothbaum BO, Foa EB** (1995). A prospective examination of symptoms of posttraumatic stress disorder in victims of nonsexual assault. *Journal of Interpersonal Violence* **10**, 201–214.
- Rothbaum BO, Foa EB, Riggs DS, Murdock T, Walsh W** (1992). A prospective examination of posttraumatic stress disorder in rape victims. *Journal of Traumatic Stress* **5**, 455–475.
- Scully JH** (2000). The American Psychiatric Association textbook of psychiatry, 3rd edition. *Journal of Clinical Psychiatry* **61**, 306–306.
- See J, MacLeod C, Bridle R** (2009). The reduction of anxiety vulnerability through the modification of attentional bias: a real-world study using a home-based cognitive bias modification procedure. *Journal of Abnormal Psychology* **118**, 65–75.
- Shane MS, Peterson JB** (2007). An evaluation of early and late stage attentional processing of positive and negative information in dysphoria. *Cognition and Emotion* **21**, 789–815.
- Shechner T, Britton JC, Perez-Edgar K, Bar-Haim Y, Ernst M, Fox NA, Leibenluft E, Pine DS** (2012). Attention biases, anxiety, and development: toward or away from threats or rewards? *Depression and Anxiety* **29**, 282–294.
- Spitzer RL, Kroenke K, Williams JBW** (1999). Validation and utility of a self-report version of PRIME-MD: the PHQ primary care study. *Journal of the American Medical Association* **282**, 1737–1744.
- Thrasher SM, Dalgleish T, Yule W** (1994). Information processing in post-traumatic stress disorder. *Behaviour Research and Therapy* **32**, 247–254.
- van der Kolk BA, van der Hart O, Marmar CR** (1996). Dissociation and information processing in posttraumatic stress disorder. In *Traumatic Stress: The Effects of Overwhelming Experience on Mind, Body, and Society* (ed. B. A. van der Kolk, A. C. McFarlane and L. Weisaeth), pp. 303–327. Guilford Press: New York.
- Wald I, Degnan KA, Gorodetsky E, Charney DS, Fox NA, Fruchter E, Goldman D, Lubin G, Pine DS, Bar-Haim Y** (2013). Attention to threats and combat-related posttraumatic stress symptoms: prospective associations and moderation by the serotonin transporter gene. *Journal of the American Medical Association Psychiatry* **70**, 401–408.
- Wald I, Lubin G, Holoshitz Y, Muller D, Fruchter E, Pine DS, Charney DS, Bar-Haim Y** (2011a). Battlefield-like stress following simulated combat and suppression of attention bias to threat. *Psychological Medicine* **41**, 699–707.
- Wald I, Shechner T, Bitton S, Holoshitz Y, Charney DS, Muller D, Fox NA, Pine DS, Bar-Haim Y** (2011b). Attention bias away from threat during life threatening danger predicts PTSD symptoms at one-year follow-up. *Depression and Anxiety* **28**, 406–411.
- Yiend J** (2010). The effects of emotion on attention: a review of attentional processing of emotional information. *Cognition and Emotion* **24**, 3–47.